# MAKING WAVES (Postprint)

**Edward W. Cliver** 

**06 October 2013** 

**Technical Paper** 

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.



AIR FORCE RESEARCH LABORATORY
Space Vehicles Directorate
3550 Aberdeen Ave SE
AIR FORCE MATERIEL COMMAND
KIRTLAND AIR FORCE BASE, NM 87117-5776

### **DTIC COPY**

## NOTICE AND SIGNATURE PAGE

Using Government drawings, specifications, or other data included in this document for any purpose other than Government procurement does not in any way obligate the U.S. Government. The fact that the Government formulated or supplied the drawings, specifications, or other data does not license the holder or any other person or corporation; or convey any rights or permission to manufacture, use, or sell any patented invention that may relate to them.

This report was cleared for public release by the 377 ABW Public Affairs Office and is available to the general public, including foreign nationals. Copies may be obtained from the Defense Technical Information Center (DTIC) (http://www.dtic.mil).

AFRL-RV-PS-TP-2014-0001 HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION IN ACCORDANCE WITH ASSIGNED DISTRIBUTION STATEMENT.

//SIGNED//

Dr. Edward W. Cliver

Edward J. Masterson, Colonel, USAF
Program Manager, AFRL/RVBXS

Chief, Battlespace Environment Division

This report is published in the interest of scientific and technical information exchange, and its publication does not constitute the Government's approval or disapproval of its ideas or findings.


this burden to Department of E 4302. Respondents should be	Defense, Washington Headquar aware that notwithstanding an	ters Services, Directorate for Info	rmation Operations and Reports ( n shall be subject to any penalty f	(0704-0188), 1215 Jeffe	erson Davis Highway, Suite 1204, Arlington, VA 22202- n a collection of information if it does not display a currently
1. REPORT DATE (DI	D-MM-YYYY)	2. REPORT TYPE			DATES COVERED (From - To)
06-10-2013		Technical Paper			1 Jul 2012 - 30 Sep 2013
4. TITLE AND SUBTIT Making Waves (Pos				5a.	CONTRACT NUMBER
				5b.	GRANT NUMBER
					PROGRAM ELEMENT NUMBER 02F
<b>6. AUTHOR(S)</b> Edward W. Cliver				230	
				PPI	<b>TASK NUMBER</b> M00005402
				EF	WORK UNIT NUMBER 004365
<b>7. PERFORMING ORC</b> Air Force Research		AND ADDRESS(ES)			PERFORMING ORGANIZATION REPORT IUMBER
Space Vehicles Dire	· ·			AF	RL-RV-PS-TP-2014-0001
3550 Aberdeen Ave					
Kirtland AFB, NM 87117-5776					
9. SPONSORING / MC	ONITORING AGENCY N	NAME(S) AND ADDRES	S(ES)		SPONSOR/MONITOR'S ACRONYM(S) RL/RVBXS
					SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited. (377ABW-2013-0759; dtd 12 Sep 2013)					
13. SUPPLEMENTARY NOTES Accepted for publication in Nature Physics, Published online: 06 October 2013. Government Purpose Rights.					
Carley, et al., (2013) link together the key elements for acceleration of particles by magnetohydrodynamic shocks at the Sun for an eruptive event on 22 September 2011. In brief, in this event the lateral expansion of a coronal mass ejection drove a quasi-perpendicular shock wave that accelerated electrons in intermittent bursts, with the intermittency attributed to the rippled or wavy surface of the shock front. The history of research on large-scale solar waves is reviewed. Such waves were discovered by radio means in 1947. Recent progress has been primarily based on high-cadence images of such waves at extreme ultra-violet wavelengths.					
15. SUBJECT TERMS					
solar activity, coronal mass ejections, large-scale waves, Moreton waves, type II radio bursts, solar energetic particles					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON E. W. Cliver
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	Unlimited	8	19b. TELEPHONE NUMBER (include area code)

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the

REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188



### **SOLAR PHYSICS**

# Making waves

High-cadence images link the phenomena required for particle acceleration at the Sun. A plasmoid-driven shock wave accelerates electrons in intermittent bursts.

### Edward W. Cliver

hock waves are a primary mechanism for the acceleration of particles at the Sun and in other astrophysical settings, such as supernovae. Shocks at the Sun and in the heliosphere offer the best chance to investigate the shock particle-acceleration process in detail. Eoin Carley and his co-workers, as they now report in Nature Physics1, have used highcadence solar images and radio spectroscopy to link together the key elements of particle acceleration at a solar shock wave: a driving plasmoid or coronal mass ejection (CME); a propagating coronal bright front at the leading edge of the CME with an associated metric radio 'type II' burst — the classic signature of a solar shock; and the coronal and interplanetary electrons accelerated at the shock. They present evidence that the shock is quasiperpendicular and attribute the bursty or quasiperiodic nature of the radio emission emanating from the type II burst to a rippled or wavy shock surface.

On 22 September 2011, a CME — an erupting bubble of magnetized plasma was observed off the limb of the Sun. The eruption was recorded by coronagraphs and extreme-ultraviolet (EUV) imagers on the Solar and Heliospheric Observatory (SOHO), the Solar Dynamics Observatory (SDO) and the Solar Terrestrial Relations Observatory (STEREO), as well as by ground-based radio telescopes. Carley et al. used coronagraph images from SOHO (near Earth) and STEREO-B (lagging ~95° behind Earth in a 1 AU orbit) to identify a faint front off the southern flank of the laterally expanding CME as a shock wave. At the same location, SDO high-cadence EUV images revealed a coronal bright front propagating southwards along the limb of the Sun in concert with an imaged type II radio source. The radio spectrogram of the event exhibited a 'herringbone' structure emanating from a nondrifting (constant frequency) backbone, suggesting electron acceleration by a shock wave moving parallel to the Sun's surface.

Taken together, these observations indicate a quasiperpendicular shock wave (propagating normal to the ambient magnetic field) driven by the lateral expansion of a CME. The shock




Figure 1 | Coronal mass ejection. The CME observed on 22 September 2011 drove a coronal bright front (CBF) and quasiperpendicular  $(Q_1)$  shock southwards along the limb of the Sun. Shock-accelerated electrons escape to interplanetary space along open field lines.

accelerates electrons, which flow inwards towards the Sun and outwards into the interplanetary medium on upstream open field lines. The electron streams produced herringbones with 2–11 s periodicities. Carley *et al.* suggest that a rippled shock surface modulated the efficiency of electron acceleration in this event, resulting in the observed time structure.

The study of large-scale waves on the Sun began in the late 1940s with the discovery of metric type II radio bursts that are characterized by a drift to lower frequencies at a rate of  $\sim 0.5$  MHz s<sup>-1</sup> (ref. 2). The causative disturbance, later identified as a magnetohydrodynamic shock wave, propagates outwards through the corona with a speed of 500-750 km s<sup>-1</sup>. The optical counterparts of type II bursts — observed in 6563-Å emission from hydrogen – propagate along the solar surface with comparable speeds3. In a key development, it was proposed that the compression of the chromosphere that is responsible for the optical wave is due to the 'sweeping skirt' of a coronal type II shock wave<sup>4</sup>. In 1998 research on large-scale solar waves was given fresh impetus when the EUV telescope on SOHO detected coronal waves<sup>5</sup>. Large-scale waves were subsequently reported in soft X-rays, 10830-Å emission from helium atoms and microwaves. A close kinematical relationship between the various manifestations of large-scale waves implied a common origin<sup>6</sup>.

In the past several years, solar scientists have used the new high-cadence imagery from the STEREO spacecraft (launched in 2006) and SDO (from 2010) to establish a strong link between CMEs and large-scale waves<sup>7</sup>. In short, CMEs drive these waves. Progress in this area includes recognition of the importance of the lateral (rather than radial) expansion of CMEs for surface wave creation8, culminating in the unambiguous identification of a quasiperpendicular type II shock wave at the Sun by Carley and team1 (Fig. 1). In addition, a clear distinction is forming between moving EUV phenomena (called pseudo waves) associated with the CME itself as a result of field line stretching

or magnetic restructuring<sup>9</sup>, and true waves that are initially driven by the CME and later become freely propagating when the lateral expansion of the CME stops<sup>10,11</sup>.

The work of Carley and colleagues also impacts research on the acceleration of solar energetic particles, or SEPs, by shock waves. At present, the surface structure of shocks, including ripples, is beyond the reach of solar imagery but, as demonstrated by Carley et al., the macro-geometry (quasiparallel or quasiperpendicular) can be determined with confidence in certain cases. The potential importance of shock orientation for SEPs was fully appreciated only recently when compositional signatures of quasiparallel and quasiperpendicular shock acceleration in large SEP events were identified<sup>12</sup>. In a recent examination of a CME, EUV wave, and SEP event on 21 March 201113, evidence from

SEP composition near-Earth measurements was reported that was consistent with the picture presented in ref. 12 although confirming radio images of a shock wave were not available. The outlook for radio imagery is promising, given the development of the Low Frequency Array in Europe and the Murchison Widefield Array in Australia. In combination with the high-cadence EUV images from STEREO and SDO, the imaging and spectroscopy capabilities of these new radio telescopes will permit more detailed study of the shock particle-acceleration process and provide a probe of the medium through which the shocks propagate<sup>14</sup>.

Edward W. Cliver is at the Space Vehicles Directorate, Air Force Research Laboratory, Sunspot, New Mexico 88349, USA. e-mail: ecliver@nso.edu

#### References

- Carley, E. P. et al. Nature Phys. http://dx.doi.org/10.1038/ nphys2767 (2013).
- Payne-Scott, R., Yabsley, D. E. & Bolton, J. G. Nature 160, 256–257 (1947).
- Moreton, G. E. & Ramsey, H. E. Publ. Astron. Soc. Pacific 72, 357 (1960).
- 4. Uchida, Y. Sol. Phys. 39, 431-449 (1974).
- 5. Thompson, B. J. et al. Geophys. Res. Lett. 25, 2465-2468 (1998).
- 6. Warmuth, A., Vršnak, B., Magdalenić, J., Hanslmeier, A. & Otruba, W. Astron. Astrophys. 418, 1117–1129 (2004).
- Veronig, A. M., Temmer, M. & Vršnak, B. Astrophys. J. 681, L113–L116 (2008).
- Kienreich, I. W., Temmer, M. & Veronig, A. M. Astrophys. J. 703, L118–L122 (2009).
- Chen, P. F., Fang, C. & Shibata, K. Astrophys. J. 622, 1202–1210 (2005).
- 10. Liu, W. et al. Astrophys. J. 753, 52 (2012).
- 11. Patsourakos, S. & Vourlidas, A. Solar Phys. 281, 187-222 (2012).
- 12. Tylka, A. J. et al. Astrophys. J. 625, 474-495 (2005).
- 13. Rouillard, A. P. et al. Astrophys. J. 752, 44 (2012).
- Gopalswamy, N., Nitta, N., Akiyama, S., Mäkelä, P. & Yashiro, S. Astrophys. J. 744, 72 (2012).

Published online: 6 October 2013

## **DISTRIBUTION LIST**

DTIC/OCP

8725 John J. Kingman Rd, Suite 0944

Ft Belvoir, VA 22060-6218 1 cy

AFRL/RVIL

Kirtland AFB, NM 87117-5776 2 cys

Official Record Copy

AFRL/RVBXS/Dr. Edward W. Cliver 1 cy

This page is intentionally left blank.